

# SEE WHAT YOU CAN ACHIEVE WITH PYZCHIVA<sup>®</sup>

(ustekinumab-ttwe)

Learn more about PYZCHIVA,  
an FDA-approved alternative  
to Stelara<sup>®</sup> (ustekinumab)



**Pyzchiva**<sup>®</sup>  
ustekinumab-ttwe | 90 mg/1 mL

## What is PYZCHIVA<sup>®</sup>?

PYZCHIVA is a prescription medicine used to treat:

- adults and children 6 years and older with moderate to severe psoriasis who may benefit from taking injections or pills (systemic therapy) or phototherapy (treatment using ultraviolet light alone or with pills)
- adults and children 6 years and older with active psoriatic arthritis
- adults 18 years and older with moderately to severely active Crohn's disease
- adults 18 years and older with moderately to severely active ulcerative colitis

## IMPORTANT SAFETY INFORMATION

### What is the most important information I should know about PYZCHIVA?

**Do not take PYZCHIVA if you are** allergic to ustekinumab or any of the ingredients in PYZCHIVA.

PYZCHIVA is a prescription medicine that affects your immune system. PYZCHIVA can increase your risk of having serious side effects, including:

**Serious Infections:** PYZCHIVA may lower your ability to fight infections and may increase your risk of infections. Some people have serious infections while taking PYZCHIVA, including tuberculosis (TB), and infections caused by bacteria, fungi, or viruses. Some people have to be hospitalized for treatment of their infection.

- Your doctor should check you for TB before starting PYZCHIVA and watch you closely for signs and symptoms of TB during treatment with PYZCHIVA

**Please see additional Important Safety Information throughout, and full Prescribing Information, including Medication Guide, for PYZCHIVA.**





**SAN DOZ**

# What you can expect with PYZCHIVA<sup>®</sup> (ustekinumab)

## What is a biosimilar?

Although they are not exactly the same, biosimilars like PYZCHIVA are similar to generic medicines. They can be used in place of name-brand drugs, delivering results that are just as safe and effective.

As a biosimilar to Stelara<sup>®</sup> (ustekinumab), PYZCHIVA offers:

- 
**An FDA-approved treatment**  
 PYZCHIVA is an FDA-approved biosimilar to Stelara<sup>®</sup>. This means PYZCHIVA has been thoroughly studied and proven to be as safe and effective as Stelara<sup>®</sup>. It can also be used in place of Stelara<sup>®</sup>.
- 
**The same symptom relief as Stelara<sup>®</sup>**  
 The FDA approved PYZCHIVA because it is proven to have the same effectiveness, safety, and quality as Stelara<sup>®</sup>. This means you can expect the same results with PYZCHIVA that you've had with Stelara<sup>®</sup>.
- 
**The same dosing schedule**  
 PYZCHIVA has the same dosing schedule as Stelara<sup>®</sup>, so switching is straightforward.
- 
**Support from day one**  
 Sandoz One Source offers comprehensive support for PYZCHIVA.



## IMPORTANT SAFETY INFORMATION (cont'd)

### What is the most important information I should know about PYZCHIVA? (cont'd)

- If your doctor feels that you are at risk for TB, you may be treated for TB before and during treatment with PYZCHIVA

You should not start taking PYZCHIVA if you have any kind of infection unless your doctor says it is okay.

#### Before starting PYZCHIVA, tell your doctor if you:

- think you have an infection or have symptoms of an infection such as:
  - fever, sweats, or chills
  - muscle aches
  - cough
  - shortness of breath
  - blood in phlegm
  - weight loss
  - warm, red, or painful skin or sores on your body
  - diarrhea or stomach pain
  - burning when you urinate or urinate more often than normal
  - feeling very tired
- are being treated for an infection or have any open cuts
- get a lot of infections or have infections that keep coming back
- have TB, or have been in close contact with someone with TB

Please see additional Important Safety Information throughout, and full Prescribing Information, including Medication Guide, for PYZCHIVA.



**Pyzchiva<sup>®</sup>**  
ustekinumab-ttwe | 90 mg/1 mL

# Enroll in the PYZCHIVA<sup>®</sup> Co-Pay Program

If you're eligible, you could pay as little as \$0 for PYZCHIVA.

- If you qualify, you can **start using your co-pay card immediately**
- Your card **doesn't expire**—you'll be automatically enrolled every year
- **Enrollment is easy**—you can enroll online or by phone

For full Terms and Conditions, visit [PYZCHIVA.com/](http://PYZCHIVA.com/)

 	
<b>Member Name:</b> <PatientFirstName> <PatientLastName>	
<b>MEDICAL CLAIMS:</b> PAYER ID: 56155 GROUP: 00003729 MEMBER ID: <XXXXXXXXXXXX>	<b>PHARMACY CLAIMS:</b> BIN: 610852 PCN: 2001 GROUP: 77770226 MEMBER ID: <XXXXXXXXXXXX>
Pay as little as \$0 per treatment	
<small>For questions, call Sandoz One Source Co-Pay Program for Pyzchiva at 1-855-SANDOZ-8 (1-855-726-3698)          Visit <a href="http://www.pyzchiva.com">www.pyzchiva.com</a> for program eligibility details.</small>	
<small>MLR-1650-US 11/2024</small>	

Whether you're starting or switching to PYZCHIVA, our team is here to help at every step



## Case Managers

- Provide education about PYZCHIVA
- Help identify opportunities for savings



## Injection support

- Offers personalized, 1:1 training options
- Nurse support to answer treatment questions



## Insurance support

- Assistance with navigating and understanding your health plan benefits
- Help with identifying a pharmacy that will arrange shipment of your medication to you



## Questions about Sandoz One Source?

Call **1-855-SANDOZ-8 (1-855-726-3698)**, or visit [PYZCHIVA.com/](http://PYZCHIVA.com/) to learn more about the Co-Pay Program and our services

## IMPORTANT SAFETY INFORMATION (cont'd)

**After starting PYZCHIVA, call your doctor right away** if you have any symptoms of an infection (see above). These may be signs of infections such as chest infections, or skin infections or shingles that could have serious complications. PYZCHIVA can make you more likely to get infections or make an infection that you have worse. People who have a genetic problem where the body does not make any of the proteins interleukin 12 (IL-12) and interleukin 23 (IL-23) are at a higher risk for certain serious infections. These infections can spread throughout the body and cause death. People who take PYZCHIVA may also be more likely to get these infections.

**Cancers:** PYZCHIVA may decrease the activity of your immune system and increase your risk for certain types of cancers. Tell your doctor if you have ever had any type of cancer. Some people who are receiving PYZCHIVA and have risk factors for skin cancer have developed certain types of skin cancers. Tell your doctor if you develop any new skin growths.

**Please see additional Important Safety Information throughout, and full Prescribing Information, including Medication Guide, for PYZCHIVA.**



ustekinumab-ttwe | 90 mg/1 mL

# Taking PYZCHIVA®

PYZCHIVA is given as a shot under the skin

The prefilled syringe injects the medicine as you manually push the plunger



If you or a caregiver will be giving your PYZCHIVA shots, it is important to be fully trained by a healthcare provider. For complete information on how to inject PYZCHIVA, please see the [Instructions for Use](#) in the [Prescribing Information](#).

**NOTE:** If you have Crohn's disease or ulcerative colitis and are not switching from Stelara®, your first dose of PYZCHIVA will be given intravenously (through an IV) before you begin your shots.

**PYZCHIVA is also available in a 45 mg vial for in-office or at-home administration by a healthcare provider.**

## IMPORTANT SAFETY INFORMATION (cont'd)

**Posterior Reversible Encephalopathy Syndrome (PRES):** PRES is a rare condition that affects the brain and can cause death. The cause of PRES is not known. If PRES is found early and treated, most people recover. Tell your doctor right away if you have any new or worsening medical problems including headache, seizures, confusion, or vision problems.

**Serious Allergic Reactions:** Serious allergic reactions can occur. Stop using PYZCHIVA and get medical help right away if you have any symptoms of a serious allergic reaction such as feeling faint, swelling of your face, eyelids, tongue, or throat, chest tightness, or skin rash.

**Lung Inflammation:** Cases of lung inflammation have happened in some people who receive PYZCHIVA and may be serious. These lung problems may need to be treated in a hospital. Tell your doctor right away if you develop shortness of breath or a cough that doesn't go away during treatment with PYZCHIVA.

Please see additional Important Safety Information throughout, and full [Prescribing Information](#), including [Medication Guide](#), for PYZCHIVA.



Pyzchiva®  
ustekinumab-ttwe | 90 mg/1 mL



# Starting or switching to PYZCHIVA®



## If you are starting treatment with PYZCHIVA:

- Your healthcare provider will talk to you about how often you'll get your PYZCHIVA shots
  - If you have Crohn's disease or ulcerative colitis, you will have a starter dose that is given intravenously (through an IV) before you begin your shots
- Your healthcare provider can give you your shots in their office, or you or a caregiver can be trained to inject your PYZCHIVA shots at home



## If you are switching to PYZCHIVA:

- PYZCHIVA has the same dosing schedule as Stelara®, so switching is straightforward
  - There should be no changes to when you take your medicine
- Your healthcare provider can give you your shots in their office, or you or a caregiver can be trained to inject your PYZCHIVA shots at home

## IMPORTANT SAFETY INFORMATION (cont'd)

**Before you receive PYZCHIVA, tell your doctor about all of your medical conditions, including if you:**


- have any of the conditions or symptoms listed above for serious infections, cancers, or PRES
- ever had an allergic reaction to PYZCHIVA or any of its ingredients. Ask your doctor if you are not sure
- have recently received or are scheduled to receive an immunization (vaccine). People who take PYZCHIVA should not receive live vaccines. Tell your doctor if anyone in your house needs a live vaccine. The viruses used in some types of live vaccines can spread to people with a weakened immune system and can cause serious problems. **You should not receive the BCG vaccine during the one year before receiving PYZCHIVA or one year after you stop receiving PYZCHIVA**
- have any new or changing lesions within psoriasis areas or on normal skin
- are receiving or have received allergy shots, especially for serious allergic reactions. Allergy shots may not work as well for you during treatment with PYZCHIVA. PYZCHIVA may also increase your risk of having an allergic reaction to an allergy shot.
- receive or have received phototherapy for your psoriasis
- are pregnant or plan to become pregnant. It is not known if PYZCHIVA can harm your unborn baby. You and your doctor should decide if you will receive PYZCHIVA
- received PYZCHIVA while you were pregnant. It is important that you tell your baby's healthcare provider before any vaccinations are given to your baby
- are breastfeeding or plan to breastfeed. PYZCHIVA can pass into your breast milk
- Talk to your doctor about the best way to feed your baby if you receive PYZCHIVA

**Tell your doctor about all the medicines you take,** including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Know the medicines you take. Keep a list of them to show your doctor and pharmacist when you get a new medicine.

**Please see additional Important Safety Information throughout, and full Prescribing Information, including Medication Guide, for PYZCHIVA.**



A man in a grey t-shirt and blue shorts is playing pickleball on a blue court. He is holding a pink paddle and a pink ball, ready to hit. The background shows a chain-link fence and green trees under a blue sky with white clouds.

# SEE WHAT YOU CAN ACHIEVE WITH PYZCHIVA<sup>®</sup>

(ustekinumab-ttwe)

Visit [PYZCHIVA.com/](https://www.pyzchiva.com/) to learn more

## IMPORTANT SAFETY INFORMATION (cont'd)

### When prescribed PYZCHIVA:

- Use PYZCHIVA exactly as your doctor tells you to
- PYZCHIVA is intended for use under the guidance and supervision of your doctor. In children 6 years and older, it is recommended that PYZCHIVA be administered by a healthcare provider. If your doctor decides that you or a caregiver may give your injections of PYZCHIVA at home, you should receive training on the right way to prepare and inject PYZCHIVA. Your doctor will determine the right dose of PYZCHIVA for you, the amount for each injection, and how often you should receive it. Do not try to inject PYZCHIVA yourself until you or your caregiver have been shown how to inject PYZCHIVA by your doctor or nurse

**Keep PYZCHIVA and all medicines out of the reach of children.**

**Common side effects for PYZCHIVA include** nasal congestion, sore throat, and runny nose; upper respiratory infections; fever; headache; tiredness; itching; nausea and vomiting; redness at the injection site; vaginal yeast infections; urinary tract infections; sinus infection; bronchitis; diarrhea; stomach pain; and joint pain. These are not all of the possible side effects with PYZCHIVA. Tell your doctor about any side effect that you experience. Ask your doctor or pharmacist for more information.

**Please see your doctor to discuss any questions you may have.**

**Please see additional Important Safety Information throughout, and full Prescribing Information, including Medication Guide, for PYZCHIVA.**

**You are encouraged to report negative side effects of prescription drugs to the FDA.**

**Visit [www.fda.gov/medwatch](https://www.fda.gov/medwatch) or call 1-800-FDA-1088.**

All trademarks and trade names are the property of their respective owners.

**SANDOZ**

© 2025 Sandoz Inc. 100 College Road West, Princeton, NJ 08540  
[www.sandoz.com](https://www.sandoz.com) MLR-1537-US 02/2025

The Pyzchiva logo features a stylized 'P' with a green and pink gradient, followed by the word 'Pyzchiva' in a blue sans-serif font. Below it, the text 'ustekinumab-ttwe | 90 mg/1 mL' is written in a smaller blue font.

**Pyzchiva<sup>®</sup>**  
ustekinumab-ttwe | 90 mg/1 mL